

Product Data Sheet

MED15 Blocking Peptide

Catalog # Source Reactivity Applications

CBP2675 Synthetic H, M, R, Z BL

Description The peptide is used to block Anti-MED15 Antibody (#CPA2675) reactivity.

Form Lyophilized powder

Gene Symbol MED15

Alternative Names ARC105; CTG7A; PCQAP; TIG1; TNRC7; Mediator of RNA polymerase II transcription

subunit 15; Activator-recruited cofactor 105 kDa component; ARC105; CTG repeat

protein 7a; Mediator complex subunit 15; Positive cofactor 2

glutamine/Q-rich-associated protein; PC2 glutamine/Q-rich-associated protein;

TPA-inducible gene 1 protein; TIG-1; Trinucleotide repeat-containing gene 7 protein

Entrez Gene 51586 (Human)

SwissProt Q9NY74 (Human)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

Directions for Use Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

Storage/Stability Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com