

## MST1/2 (pT183) Blocking Peptide

Catalog #	Source	Reactivity	Applications
CBP4262	Synthetic	H, M, R, B, C, P, Z	BL
<b>Description</b>	The peptide is used to block Anti-MST1/2 (pT183) Antibody (#CPA4262) reactivity.		
<b>Form</b>	Lyophilized powder		
<b>Gene Symbol</b>	STK3; STK4		
<b>Alternative Names</b>	STK3; KRS1; MST2; Serine/threonine-protein kinase 3; Mammalian STE20-like protein kinase 2; MST-2; STE20-like kinase MST2; Serine/threonine-protein kinase Krs-1; STK4; KRS2; MST1; Serine/threonine-protein kinase 4; Mammalian STE20-like protein kinase 1; M		
<b>Entrez Gene</b>	6789 (Human); 56274, 58231 (Mouse); 65189 (Rat)		
<b>SwissProt</b>	Q13188, Q13043 (Human); Q9JI10, Q9JI11 (Mouse); O54748 (Rat)		
<b>Purity</b>	>85%		
<b>Quality Control</b>	The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry.		
<b>Directions for Use</b>	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours.		
<b>Storage/Stability</b>	Shipped at 4°C. Store at -20°C for one year.		

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**  
[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**  
[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**  
[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**  
[custom@cohesionbio.com](mailto:custom@cohesionbio.com)