

Product Data Sheet

CD147 Blocking Peptide

Catalog # Source Reactivity Applications

CBP5269 Synthetic H, R, B BL

Description The peptide is used to block Anti-CD147 Antibody (#CPA5269) reactivity.

Form Lyophilized powder

Gene Symbol BSG

Alternative Names Basigin; 5F7; Collagenase stimulatory factor; Extracellular matrix metalloproteinase

inducer; EMMPRIN; Leukocyte activation antigen M6; OK blood group antigen;

Tumor cell-derived collagenase stimulatory factor; TCSF; CD147

Entrez Gene 682 (Human); 25246 (Rat)

SwissProt P35613 (Human); P26453 (Rat)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

Directions for UseBlocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

Storage/Stability Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED