

Goat Anti-Mouse IgG (H&L) TRITC

Catalog #	Source	Reactivity	Applications
CSA1022	Goat	M	IF, FC
Description	Goat Polyclonal Secondary Antibody to Mouse IgG (H&L) TRITC		
Immunogen	Mouse IgG (H&L)		
Purification	The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.		
Specificity	By immunoelectrophoresis and ELISA this antibody reacts specifically with Mouse IgG. No antibody was detected against non immunoglobulin serum proteins.		
Clonality	Polyclonal		
Conjugation	TRITC (Rhodamine)		
Form	Liquid in 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, containing 1% BSA, 50% glycerol, 0.05% Sodium Azide		
Dilution	IF (1/50 - 1/200), FC (1/50 - 1/200)		
Storage/Stability	Shipped at 4°C. Upon delivery aliquot and store at -20°C for six months. Avoid freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com