

Storage/Stability

Product Data Sheet

Goat Anti-Rabbit IgG (H&L) TRITC

freeze/thaw cycles.

Source	Reactivity	Applications	
Goat	Rb	IF, FC	
Goat	Goat Polyclonal Secondary Antibody to Rabbit IgG (H&L) TRITC		
Rabb	Rabbit IgG (H&L)		
The a	The antibody was isolated from antisera by immunoaffinity chromatography using		
antig	antigens coupled to agarose beads.		
By im	By immunoelectrophoresis and ELISA this antibody reacts specifically with Rabbit		
IgG. I	No antibody was dete	cted against non immunoglobulin serum proteins.	
Polyc	lonal		
TRITO	TRITC (Rhodamine)		
Liqui	Liquid in 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, containing 1% BSA, 50%		
glyce	rol, 0.05% Sodium Az	ide	
IF (1/5	50 - 1/200), FC (1/50 -	1/200)	
	Goat Rabb The a antig By im IgG. N Polyc TRITC Liquid glyce	Goat Polyclonal Secondary Rabbit IgG (H&L) The antibody was isolated antigens coupled to agaros By immunoelectrophoresis IgG. No antibody was determined to the polyclonal TRITC (Rhodamine)	

Shipped at 4°C. Upon delivery aliquot and store at -20°C for six months. Avoid

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com