

Mouse Anti-Monkey IgG (H&L) Biotin

Catalog #	Source	Reactivity	Applications
CSA2240	Mouse	Mk	E, WB, IH, IC
Description	Mouse Monoclonal Secondary Antibody to Monkey IgG (H&L) Biotin		
Immunogen	Monkey IgG		
Purification	The antibody was isolated from ascitic by immunoaffinity chromatography using antigens coupled to agarose beads.		
Specificity	By immunoelectrophoresis and ELISA this antibody reacts specifically with Monkey IgG. Do not cross reactivity with human IgG. No antibody was detected against non immunoglobulin serum proteins.		
Clonality	Monoclonal		
Conjugation	Biotin		
Form	Liquid in 0.01M Phosphate Buffered Saline, pH 7.2, containing 1% BSA, 50% glycerol, 0.05% Sodium Azide		
Dilution	E (1/5000 - 1/20000), WB (1/5000 - 1/20000), IH (1/500 - 1/1000), IC (1/500 - 1/1000)		
Storage/Stability	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com