

Product Data Sheet

Goat Anti-Mouse IgA - FITC

Catalog #	Source	e Reactivity	Applications	
CSA2307	Goat	М	E, IH, IF, FC	
Description		Goat Polyclonal Secondary Antibody to Mouse IgA FITC		
Immunogen		Mouse IgA		
Purification		The antibody was isolated	from antisera by immunoaffinity chromatography using	
		antigens coupled to agaros	se beads.	
Specificity By immu		By immunoelectrophoresis	unoelectrophoresis and ELISA this antibody reacts specifically with Mouse	
		IgA. No antibody was dete	cted against non immunoglobulin serum proteins.	
Clonality		Polyclonal		
Conjugation		FITC		
Form		1 mg/ml. Liquid in 0.01M F	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,	
		50% glycerol, 0.05% Sodiu	n Azide	
Dilution		IH (1/100 - 1/1000), IF (1/10	0 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay	
		dependent concentration)		
Fluorophore		Ex/Em = 493/528 nm		
Storage/Stabi	lity	Shipped at 4°C. Upon deliv	ery aliquot and store at -20°C for one year. Avoid	
		freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com