

Mouse Anti-Human IgG Fc - FITC

Catalog #	Source	Reactivity	Applications
CSA2338	Mouse	H	E, IH, IF, FC
Description	Mouse Monoclonal Secondary Antibody to Human IgG Fc FITC		
Immunogen	Human IgG Fc		
Purification	The antibody was isolated from ascitic by immunoaffinity chromatography using antigens coupled to agarose beads.		
Specificity	By immunoelectrophoresis and ELISA this antibody reacts specifically with Human IgG Fc. No antibody was detected against non immunoglobulin serum proteins.		
Clonality	Monoclonal		
Conjugation	FITC		
Form	1 mg/ml. Liquid in 0.01M Phosphate Buffered Saline, pH 7.2, containing 1% BSA, 50% glycerol, 0.05% Sodium Azide		
Dilution	IH (1/100 - 1/1000), IF (1/100 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay dependent concentration)		
Fluorophore	Ex/Em = 493/528 nm		
Storage/Stability	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com