

Product Data Sheet

CD168 Blocking Peptide

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|------------|---|---|--|--|--|
| CBP1532 | Synthetic | H, M, R | BL | | | |
| Description | The p | The peptide is used to block Anti-CD168 Antibody (#CPA1532) reactivity. | | | | |
| Form | Lyoph | ilized powder | | | | |
| Gene Symbol | HMM | HMMR | | | | |
| Alternative N | ames IHABP | IHABP; RHAMM; Hyaluronan mediated motility receptor; Intracellular hyaluronic | | | | |
| | acid-b | inding protein; Rece | ptor for hyaluronan-mediated motility; CD168 | | | |
| Entrez Gene | 3161 | 3161 (Human); 15366 (Mouse) | | | | |
| SwissProt | 07533 | O75330 (Human); Q00547 (Mouse); P97779 (Rat) | | | | |
| Purity | >85% | | | | | |
| Quality Contr | ol The qu | The quality of the peptide was evaluated by reversed-phase HPLC and by mass | | | | |
| | spectr | rometry. | | | | |
| Directions for | Use Blocki | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to | | | | |
| | antibo | ody) and incubate the | e mixture at 4°C for overnight or at room temperature for | | | |
| | 2 hou | rs. | | | | |
| Storage/Stabi | lity Shipp | ed at 4°C. Store at -2 | 0°C for one year. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |