

Product Data Sheet

ERK1/2 Blocking Peptide

Catalog # Source Reactivity Applications

CBP1944 Synthetic H, M, R, B, D, P BL

Description The peptide is used to block Anti-ERK1/2 Antibody (#CPA1944) reactivity.

Form Lyophilized powder

Gene Symbol MAPK1; MAPK3

Alternative Names MAPK3; ERK1; PRKM3; Mitogen-activated protein kinase 3; MAP kinase 3; MAPK 3;

ERT2; Extracellular signal-regulated kinase 1; ERK-1; Insulin-stimulated MAP2 kinase;

MAP kinase isoform p44; p44-MAPK; Microtubule-associated protein 2 kinase;

p44-ERK1; MAPK1; ERK2; PRKM1; PRKM2; Mitogen-activated protein kinase 1; MAP

kinase 1; MAPK 1; ERT1; Extracellular signal-regulated kinase 2; ERK-2; MAP kinase

isoform p42; p42-MAPK; Mitogen-activated protein kinase 2; MAP kinase 2; MAPK 2

Entrez Gene 5595 (Human); 26417, 26413 (Mouse); 50689, 116590 (Rat)

SwissProt P27361, P28482 (Human); Q63844, P63085 (Mouse); P21708, P63086 (Rat)

Purity >85%

Quality Control

The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

Directions for Use Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

Storage/Stability Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com