

Product Data Sheet

RAC2 Blocking Peptide

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|------------|---|---|--|--|--|
| CBP3002 | Synthetic | H, M, R | BL | | | |
| Description | The J | peptide is used to bloc | k Anti-RAC2 Antibody (#CPA3002) reactivity. | | | |
| Form | Lyop | Lyophilized powder | | | | |
| Gene Symbol | RAC2 | RAC2 | | | | |
| Alternative N | ames Ras-r | Ras-related C3 botulinum toxin substrate 2; GX; Small G protein; p21-Rac2 | | | | |
| Entrez Gene | 5880 | 5880 (Human); 19354 (Mouse) | | | | |
| SwissProt | P151 | P15153 (Human); Q05144 (Mouse) | | | | |
| Purity | >85% | >85% | | | | |
| Quality Contr | ol The o | The quality of the peptide was evaluated by reversed-phase HPLC and by mass | | | | |
| | spec | trometry. | | | | |
| Directions for | Use Block | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to | | | | |
| | antik | oody) and incubate the | e mixture at 4°C for overnight or at room temperature for | | | |
| | 2 ho | urs. | | | | |
| Storage/Stab | ility Ship | oed at 4°C. Store at -20 |)°C for one year. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |