

Product Data Sheet

ERK2 Blocking Peptide

| Catalog # | Source | Reactivity | Applications |
|---------------------------|--|---------------------|--------------|
| CBP3298 | Synthetic | H, M, R, B, C, P, Z | BL |
| Description | The peptide is used to block Anti-ERK2 Antibody (#CPA3298) reactivity. | | |
| Form | Lyophilized powder | | |
| Gene Symbol | MAPK1 | | |
| Alternative Names | ERK2; PRKM1; PRKM2; Mitogen-activated protein kinase 1; MAP kinase 1; MAPK 1; ERT1; Extracellular signal-regulated kinase 2; ERK-2; MAP kinase isoform p42; p42-MAPK; Mitogen-activated protein kinase 2; MAP kinase 2; MAPK 2 | | |
| Entrez Gene | 5594 (Human); 26413 (Mouse); 116590 (Rat) | | |
| SwissProt | P28482 (Human); P63085 (Mouse); P63086 (Rat) | | |
| Purity | >85% | | |
| Quality Control | The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry. | | |
| Directions for Use | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours. | | |
| Storage/Stability | Shipped at 4°C. Store at -20°C for one year. | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com