

Product Data Sheet

MARK Blocking Peptide

Catalog #	Source	Reactivity	Applications			
CBP3486	Synthetic	H, M, R, Z	BL			
Description	The p	The peptide is used to block Anti-MARK Antibody (#CPA3486) reactivity.				
Form	Lyoph	Lyophilized powder				
Gene Symbol	MARK	MARK1; MARK2; MARK3; MARK4				
Alternative N	ames MARk	MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1;				
	MAP/	MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c;				
	MARK	MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1;				
	MAP/	microtubule affinity-r	regulating			
Entrez Gene	4139;	4139; 2011; 4140; 57787 (Human); 226778; 13728; 17169; 232944 (Mouse);				
	11701	l6; 60328; 170577 (Ra	at)			
SwissProt	Q9P0	Q9P0L2; Q7KZI7; P27448; Q96L34 (Human); Q8VHJ5; Q05512; Q03141; Q8CIP4				
	(Mou	se); 008678; 008679;	; Q8VHF0 (Rat)			
Purity	>85%	>85%				
Quality Contr	ol The q	The quality of the peptide was evaluated by reversed-phase HPLC and by mass				
	spect	rometry.				
Directions for	Use Blocki	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to				
	antibo	ody) and incubate the	e mixture at 4°C for overnight or at room temperature for	or		
	2 hou	rs.				
Storage/Stabi	i lity Shipp	ed at 4°C. Store at -20)°C for one year.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com