

Product Data Sheet

Histone Deacetylase 8 Blocking Peptide

Catalog #	Source	Reactivity	Applications			
CBP3992	Synthetic	H, M, R, Mk	BL			
Description	The p	eptide is used to blocl	k Anti-Histone Deacetylase 8 Antibody (#CPA3992)		
	reacti	vity.				
Form	Lyoph	ilized powder				
Gene Symbol	HDAC	HDAC8				
Alternative N	ames HDAC	HDACL1; Histone deacetylase 8; HD8				
Entrez Gene	55869	55869 (Human); 70315 (Mouse); 100911968, 363481 (Rat)				
SwissProt	Q9BY	Q9BY41 (Human); Q8VH37 (Mouse); B1WC68 (Rat)				
Purity	>85%					
Quality Contr	ol The q	The quality of the peptide was evaluated by reversed-phase HPLC and by mass				
	spect	rometry.				
Directions for	Use Block	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to				
	antibo	ody) and incubate the	mixture at 4°C for overnight or at room	temperature for		
	2 hou	rs.				
Storage/Stab	i lity Shipp	ed at 4°C. Store at -20)°C for one year.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com