

Product Data Sheet

TAK1 Blocking Peptide

Catalog # Source Reactivity Applications

CBP4075 Synthetic H, M, R, B, P, Z BL

Description The peptide is used to block Anti-TAK1 Antibody (#CPA4075) reactivity.

Form Lyophilized powder

Gene Symbol MAP3K7

Alternative Names TAK1; Mitogen-activated protein kinase kinase kinase 7; Transforming growth

factor-beta-activated kinase 1; TGF-beta-activated kinase 1

Entrez Gene 6885 (Human); 26409 (Mouse); 100910771, 313121 (Rat)

SwissProt 043318 (Human); Q62073 (Mouse); P0C8E4 (Rat)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

Directions for Use Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

Storage/Stability Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED