

Product Data Sheet

NFAT4 (pS165) Blocking Peptide

Catalog # Source Reactivity Applications

CBP4119 Synthetic H, M, Mk BL

Description The peptide is used to block Anti-NFAT4 (pS165) Antibody (#CPA4119) reactivity.

Form Lyophilized powder

Gene Symbol NFATC3

Alternative Names NFAT4; Nuclear factor of activated T-cells, cytoplasmic 3; NF-ATc3; NFATc3; NFATx;

T-cell transcription factor NFAT4; NF-AT4

Entrez Gene 4775 (Human)

SwissProt Q12968 (Human); P97305 (Mouse)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

Directions for UseBlocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

Storage/Stability Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com