

Product Data Sheet

PKC mu (pY463) Blocking Peptide

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|-------------------|---|---|----|--|--|
| CBP4171 | Synthetic | H, M, R, D, Mk | BL | | | |
| Description | The p | The peptide is used to block Anti-PKC mu (pY463) Antibody (#CPA4171) reactivity. | | | | |
| Form | Lyoph | Lyophilized powder | | | | |
| Gene Symbol | PRKD | PRKD1 | | | | |
| Alternative N | ames PKD; F | PKD; PKD1; PRKCM; Serine/threonine-protein kinase D1; Protein kinase C mu type; | | | | |
| | Protei | n kinase D; nPKC-D1; n | nPKC-mu | | | |
| Entrez Gene | 5587 (| 5587 (Human); 18760 (Mouse); 85421 (Rat) | | | | |
| SwissProt | Q1513 | Q15139 (Human); Q62101 (Mouse); Q9WTQ1 (Rat) | | | | |
| Purity | >85% | >85% | | | | |
| Quality Contr | ol The qu | The quality of the peptide was evaluated by reversed-phase HPLC and by mass | | | | |
| | spectr | ometry. | | | | |
| Directions for | Use Blocki | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to | | | | |
| | antibo | ody) and incubate the n | mixture at 4°C for overnight or at room temperature for | or | | |
| | 2 hou | rs. | | | | |
| Storage/Stab | ility Shippe | ed at 4°C. Store at -20° | C for one year. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |