

Product Data Sheet

Histone H3 Blocking Peptide

Catalog #	Source	Reactivity	Applications			
CBP7033	Synthetic	H, M, R, B, C, P, Z	BL			
Description	The	peptide is used to block A	nti-Histone H3 Antibody (#CPA7033) reactivity.			
Form	Lyop	hilized powder				
Gene Symbol	HIST	HIST1H3A				
Alternative Na	ames HIST	HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E;				
	H3F	H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF;				
	HIST	HIST1H3J; H3FJ; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone				
	H3/c	; Histone H3/f; Histone H3	/			
Entrez Gene	8350	8350; 8351; 8352; 8353; 8354; 8355; 8356; 8357; 8358; 8968 (Human); 319152;				
	1507	7; 15078 (Mouse); 29115	9; 100361558 (Rat)			
SwissProt	P684	P68431; Q71DI3; P84243 (Human); P68433; P84228; P84244 (Mouse); Q6LED0;				
	P842	245 (Rat)				
Purity	>859	>85%				
Quality Contro	ol The	The quality of the peptide was evaluated by reversed-phase HPLC and by mass				
	spec	trometry.				
Directions for	Use Bloc	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to				
	antil	oody) and incubate the mi	xture at 4°C for overnight or at room temperature for			
	2 ho	urs.				
Storage/Stabi	lity Ship	oed at 4°C. Store at -20°C	for one year.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com