

Myc-tag Blocking Peptide

Catalog #	Source	Reactivity	Applications
CBP9004	Synthetic	N/A	BL

Description	The peptide is used to block Anti-Myc-tag Antibody reactivity.
Form	Lyophilized powder
Gene Symbol	
Alternative Names	c Myc; c Myc Epitope Tag; c Myc Tag; Myc Epitope Tag; Myc proto-oncogene protein; Transcription factor p64
Entrez Gene	
SwissProt	
Purity	>85%
Quality Control	The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry.
Directions for Use	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours.
Storage/Stability	Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com