

## **Product Data Sheet**

## hsa-miR-4687-5p miRNA Antagomir

Catalog #	Source	Reactivity	Applications	
CIJ1508	Synthetic	Н		
Description	<b>Description</b> Synthetic miRNA Antagomir is used to inhibit the activity of target hsa-m		nir is used to inhibit the activity of target hsa-miR-4687-5p	
	mRN	٩.		
Specificity	Antagomir is chemically-modified single-strand miRNA inhibitor functioning by			
	block	blocking miRNA regulation of target gene expression efficiently. They are synthesized		
	to red	to reduce the ability of endogenous miRNAs to silence target mRNA transcripts.		
	They	They can down-regulate the corresponding endogenous miRNAs. Our miRNA		
	antag	omir is single-strand	miRNA inhibitor carrying the chemically modifications	
	funct	ioning by blocking m	iRNA regulation of target gene expression efficiently.	
Form	Lyoph	Lyophilized powder		
Gene Symbol	hsa-n	hsa-miR-4687-5p		
Accession No.	MIMA	MIMAT0019774		
Components	This s	This synthetic miRNA is based on the mature miRNA sequence. The strand of the		

Components

This synthetic miRNA is based on the mature miRNA sequence. The strand of the antagomir has 2 phosphorothioates at the 5' end, 4 phosphorothioates, 1 cholesterol group at the 3' end, and full-length nucleotide 2'-methoxy modification. Stability of miRNA antagomir appears to be significantly higher than miRNA inhibitors. It exhibits enhanced cellular uptake, stability and regulatory activity and is recommended for miRNA functional studies in vitro and in vivo.

**Directions for Use** 

Briefly centrifuge tubes containing miRNA antagomir to ensure that the miRNA pellet is located at the bottom of the tube. Dissolve miRNA antagomir to a convenient stock concentration using the recommended volume of DEPC H2O (or RNase-free water). For example: dissolve 10 nmol miRNA antagomir to 20 μM using

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**





## **Product Data Sheet**

500 µl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

**COHESION BIOSCIENCES LIMITED**