

Product Data Sheet

mmu-miR-28a-5p miRNA Antagomir

| Catalog # | Source | Reactiv | ty Applications | | |
|----------------|---|--|--|--|--|
| CIN0163 | Syntheti | c M | | | |
| Description | S | Synthetic miRNA Antagomir is used to inhibit the activity of target mmu-miR-28a-5p | | | |
| | n | nRNA. | | | |
| Specificity | A | Antagomir is chemically-modified single-strand miRNA inhibitor functioning by | | | |
| | b | locking miRNA | regulation of target gene expression efficiently. They are synthesized | | |
| | to | to reduce the ability of endogenous miRNAs to silence target mRNA transcripts. | | | |
| | Т | hey can down-r | egulate the corresponding endogenous miRNAs. Our miRNA | | |
| | а | antagomir is single-strand miRNA inhibitor carrying the chemically mod | | | |
| | fu | unctioning by b | ocking miRNA regulation of target gene expression efficiently. | | |
| Form | Ŀ | yophilized powe | der | | |
| Gene Symbol | n | nmu-miR-28a-5 | p | | |
| Accession No. | . N | /IMAT0000653 | | | |
| Components | Т | This synthetic miRNA is based on the mature miRNA sequence. The strand of the | | | |
| | а | ntagomir has 2 | phosphorothioates at the 5' end, 4 phosphorothioates, 1 | | |
| | C | holesterol grou | o at the 3' end, and full-length nucleotide 2'-methoxy modification. | | |
| | S | tability of miRN | A antagomir appears to be significantly higher than miRNA | | |
| | ir | nhibitors. It ex | hibits enhanced cellular uptake, stability and regulatory activity and | | |
| | is | s recommended | for miRNA functional studies in vitro and in vivo. | | |
| Directions for | ctions for Use Briefly centrifuge tubes containing miRNA antagomir to ensure that the miRNA | | e tubes containing miRNA antagomir to ensure that the miRNA | | |
| | р | ellet is located | at the bottom of the tube. Dissolve miRNA antagomir to a | | |
| | C | onvenient stock | concentration using the recommended volume of DEPC H2O (or | | |
| | R | Nase-free wate | r). For example: dissolve 10 nmol miRNA antagomir to 20 μM using | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com