

# **Product Data Sheet**

### rno-miR-338-5p miRNA Antagomir

| Catalog #                 | Source   | Reactivit  | y Applications  |  |  |
|---------------------------|----------|--|---|--|--|
| CIS0236                   | Syntheti | c R  |   |  |  |
| Description               | S        | ynthetic miRNA /   | Antagomir is used to inhibit the activity of target rno-miR-338-5p    |  |  |
|                           | n        | nRNA.  |   |  |  |
| Specificity               | А        | ntagomir is chen   | nically-modified single-strand miRNA inhibitor functioning by         |  |  |
|                           | b        | locking miRNA re   | egulation of target gene expression efficiently. They are synthesized |  |  |
|                           | to       | o reduce the abil  | ity of endogenous miRNAs to silence target mRNA transcripts.          |  |  |
|                           | Т        | hey can down-re  | gulate the corresponding endogenous miRNAs. Our miRNA                 |  |  |
|                           | а        | ntagomir is single   | e-strand miRNA inhibitor carrying the chemically modifications        |  |  |
|                           | fu       | unctioning by blo  | cking miRNA regulation of target gene expression efficiently.         |  |  |
| Form                      | L        | yophilized powde   | er  |  |  |
| Gene Symbol               | rı       | no-miR-338-5p  |   |  |  |
| Accession No.             | . N      | /IMAT0004646   |   |  |  |
| Components                | Т        | his synthetic miR  | NA is based on the mature miRNA sequence. The strand of the           |  |  |
|                           | а        | ntagomir has 2 p   | hosphorothioates at the 5' end, 4 phosphorothioates, 1                |  |  |
|                           |          | cholesterol group at the 3' end, and full-length nucleotide 2'-methoxy modification. |   |  |  |
|                           |          | Stability of miRNA antagomir appears to be significantly higher than miRNA           |   |  |  |
|                           | ir       | nhibitors. It exh  | ibits enhanced cellular uptake, stability and regulatory activity and |  |  |
|                           | is       | s recommended f  | or miRNA functional studies in vitro and in vivo.                     |  |  |
| <b>Directions for Use</b> |          | Briefly centrifuge tubes containing miRNA antagomir to ensure that the miRNA         |   |  |  |
|                           | р        | ellet is located at  | t the bottom of the tube. Dissolve miRNA antagomir to a               |  |  |
|                           | C        | onvenient stock (  | concentration using the recommended volume of DEPC H2O (or            |  |  |
|                           | R        | Nase-free water)   | ). For example: dissolve 10 nmol miRNA antagomir to 20 $\mu M$ using  |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com