

Product Data Sheet

rno-miR-29b-5p miRNA Antagomir

| Catalog # | Source | Reactivity | Applications | | |
|----------------|-----------|--|--|--|--|
| CIS0254 | Synthetic | R | | | |
| Description | Sy | nthetic miRNA A | ntagomir is used to inhibit the activity of target rno-miR-29b-5p | | |
| | m | RNA. | | | |
| Specificity | A | ntagomir is chemi | cally-modified single-strand miRNA inhibitor functioning by | | |
| | bl | ocking miRNA reg | gulation of target gene expression efficiently. They are synthesized | | |
| | tc | reduce the abilit | y of endogenous miRNAs to silence target mRNA transcripts. | | |
| | Tł | ney can down-reg | ulate the corresponding endogenous miRNAs. Our miRNA | | |
| | ar | ntagomir is single- | strand miRNA inhibitor carrying the chemically modifications | | |
| | fu | nctioning by bloc | king miRNA regulation of target gene expression efficiently. | | |
| Form | Ly | ophilized powder | | | |
| Gene Symbol | rr | o-miR-29b-5p | | | |
| Accession No. | . N | IMAT0004717 | | | |
| Components | Tł | nis synthetic miRN | IA is based on the mature miRNA sequence. The strand of the | | |
| | ar | ntagomir has 2 ph | osphorothioates at the 5' end, 4 phosphorothioates, 1 | | |
| | | cholesterol group at the 3' end, and full-length nucleotide 2'-methoxy modification. | | | |
| | St | ability of miRNA a | antagomir appears to be significantly higher than miRNA | | |
| | in | hibitors. It exhil | bits enhanced cellular uptake, stability and regulatory activity and | | |
| | is | recommended fo | r miRNA functional studies in vitro and in vivo. | | |
| Directions for | Use Bi | iefly centrifuge tu | ubes containing miRNA antagomir to ensure that the miRNA | | |
| | pe | ellet is located at t | the bottom of the tube. Dissolve miRNA antagomir to a | | |
| | СС | onvenient stock co | oncentration using the recommended volume of DEPC H2O (or | | |
| | RI | Nase-free water). | For example: dissolve 10 nmol miRNA antagomir to 20 μM using | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com