

Product Data Sheet

Anti-GPRC5D Antibody

Catalog #	Source	Reactivity	Applications		
CPA5483	Rabbit	H, M, R	WB, IF/IC		
Description	Rabb	Rabbit polyclonal antibody to GPRC5D			
Immunogen	KLH-	conjugated synthetic pe	ptide encompassing a sequence within the C-term		
	regio	on of human GPRC5D. Tl	ne exact sequence is proprietary.		
Purification	The a	antibody was purified b	y immunogen affinity chromatography.		
Specificity	Reco	gnizes endogenous leve	els of GPRC5D protein.		
Clonality	Polyc	lonal			
Conjugation					
Form	Liqui	d in 0.42% Potassium p	hosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
	and (0.01% sodium azide.			
Dilution	WB (1/500 - 1/1000), IF/IC (1/	100 - 1/500)		
Gene Symbol	GPR	C5D			
Alternative Na	ames G-pro	otein coupled receptor	family C group 5 member D		
Entrez Gene	5550	7 (Human); 93746 (Moi	use)		
SwissProt	Q9N	ZD1 (Human); Q9JIL6 (N	1ouse)		
Storage/Stabi	lity Shipp	oed at 4°C. Upon delive	ry aliquot and store at -20°C for one year. Avoid		
	freez	e/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



KDa <u>A</u> <u>B</u> <u>C</u> <u>D</u> <u>E</u> 72 55 43 34 26 For research purposes only, not for human use

Product Data Sheet

Western blot analysis of GPRC5D expression in MCF7 (A), SGC7901 (B), Panc1 (C), CT26 (D), PC12 (E) whole cell lysates. (Predicted band size: 38 kD; Observed band size: 39 kD)



Immunofluorescent analysis of GPRC5D staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com