

Product Data Sheet

Anti-HIF1 beta Antibody

Catalog #	Source	Reactivity	Applications	
CPA9172	Mouse	M	WB, IH	
Description	Ν	Nouse monoclonal antiboo	ly to HIF1 beta	
Immunogen	F	Recombinant protein corre	sponding to mouse HIF1 beta.	
Purification				
Specificity	F	Recognizes endogenous lev	els of HIF1 beta protein.	
Clonality	Ν	Monoclonal		
Conjugation				
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
	а	and 0.01% sodium azide.		
Dilution	V	NB (1/1000 - 1/2000), IH (1/	100 - 1/200)	
Gene Symbol	A	ARNT		
Alternative Names		BHLHE2; Aryl hydrocarbon receptor nuclear translocator; ARNT protein; Class E basic		
	h	nelix-loop-helix protein 2; t	HLHe2; Dioxin receptor nuclear translocator;	
	ŀ	lypoxia-inducible factor 1-	beta; HIF-1-beta; HIF1-beta	
Entrez Gene	1	1863 (Mouse)		
SwissProt	P	253762 (Mouse)		
Storage/Stabi	l <mark>ity</mark> S	hipped at 4°C. Upon delive	ery aliquot and store at -20°C for one year. Avoid	
	f	reeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

Coherion

kDa A


200

140

For research purposes only, not for human use

Product Data Sheet

Western blot analysis of HIF1 beta expression in mouse brain (A) whole cell lysates. (Predicted band size: 86 kD; Observed band size: 86 kD)



Immunohistochemical analysis of HIF1 beta staining in mouse brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com