

Product Data Sheet

Anti-HSC70 Antibody

Catalog #	Source	Reactivity	Applications			
CPA9190	Mouse	H, M, R	WB, IH			
Description	Мо	use monoclonal antibo	dy to HSC70			
Immunogen	Rec	ombinant protein corre	esponding to human HSC70.			
Purification						
Specificity	Rec	Recognizes endogenous levels of HSC70 protein.				
Clonality	Мо	noclonal				
Conjugation						
Form	Liqu	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,				
	and	d 0.01% sodium azide.				
Dilution	WB	6 (1/2000 - 1/5000), IH (1	/50 - 1/200)			
Gene Symbol	HSF	YA8				
Alternative Na	ames HSC	270; HSP73; HSPA10; He	eat shock cognate 71 kDa protein; Heat shock 70 kDa			
	pro	tein 8; Lipopolysacchar	ide-associated protein 1; LAP-1; LPS-associated protein 1			
Entrez Gene	331	2 (Human)				
SwissProt	P11	142 (Human)				
Storage/Stabi	lity Shij	pped at 4°C. Upon deliv	ery aliquot and store at -20°C for one year. Avoid			
	free	eze/thaw cycles.				

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




For research purposes only, not for human use

Product Data Sheet



Western blot analysis of HSC70 expression in Hela (A), mouse brain (B), rat brain (C) whole cell lysates. (Predicted band size: 70 kD; Observed band size: 70 kD)



Immunohistochemical analysis of HSC70 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com