

Product Data Sheet

Anti-Bestrophin-1 Antibody


Catalog #	Source	Reactivity	Applications		
CPA9240	Rabbit	H, M, R	WB, IH		
Description	Rabb	Rabbit polyclonal antibody to Bestrophin-1			
Immunogen	KLH-	conjugated synthetic pept	ide encompassing a sequence of human Bestrophin-1.		
	The	exact sequence is propriet	ary.		
Purification	The	antibody was purified by in	nmunogen affinity chromatography.		
Specificity	Reco	gnizes endogenous levels	of Bestrophin-1 protein.		
Clonality	Poly	clonal			
Conjugation					
Form	Liqui	id in 0.42% Potassium pho	sphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
	and	0.01% sodium azide.			
Dilution	WB	(1/1000 - 1/2000), IH (1/50 -	1/200)		
Gene Symbol	BEST	1			
Alternative Na	ames VMD	02; Bestrophin-1; TU15B; V	itelliform macular dystrophy protein 2		
Entrez Gene	7439) (Human)			
SwissProt	0760	090 (Human)			
Storage/Stabi	lity Ship	ped at 4°C. Upon delivery	aliquot and store at -20°C for one year. Avoid		
	freez	ze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com


Coherion



For research purposes only, not for human use

Product Data Sheet

Western blot analysis of Bestrophin-1 expression in PC3 (A), mouse brain (B), rat brain (C) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 67 kD)



Immunohistochemical analysis of Bestrophin-1 staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com