

Product Data Sheet

Anti-Histone H2A.Z Antibody

Catalog #	Source	Reactivity	Applications		
CPA9325	Rabbit	M, R	WB, IH		
Description	Ra	Rabbit polyclonal antibody to Histone H2A.Z			
Immunogen	KL	LH-conjugated synthetic p	eptide encompassing a sequence of mouse Histone		
	H	2A.Z. The exact sequence	is proprietary.		
Purification	Tł	ne antibody was purified	by immunogen affinity chromatography.		
Specificity	Re	ecognizes endogenous lev	els of Histone H2A.Z protein.		
Clonality	Pc	olyclonal			
Conjugation					
Form	Lie	quid in 0.42% Potassium	ohosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
	ar	nd 0.01% sodium azide.			
Dilution	W	/B (1/1000 - 1/2000), IH (1,	200 - 1/500)		
Gene Symbol	H2	2AFZ			
Alternative Na	ames H2	2AZ; Histone H2A.Z; H2A,	z		
Entrez Gene	51	1788 (Mouse); 58940 (Ra	:)		
SwissProt	PC	0C0S6 (Mouse); P0C0S7 (Rat)		
Storage/Stabi	lity Sh	nipped at 4°C. Upon deliv	ery aliquot and store at -20°C for one year. Avoid		
	fre	eeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



For research purposes only, not for human use

Product Data Sheet

Western blot analysis of Histone H2A.Z expression in rat brain (A), mouse brain (B) whole cell lysates. (Predicted band size: 13 kD; Observed band size: 15 kD)



Immunohistochemical analysis of Histone H2A.Z staining in mouse spleen formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com