

Product Data Sheet

Anti-S100-A4 Antibody

Catalog # Source Reactivity Applications

CPA9716 Mouse H IH

Description Mouse monoclonal antibody to S100-A4

Immunogen KLH-conjugated synthetic peptide encompassing a sequence within human S100-A4.

The exact sequence is proprietary.

Purification The antibody was purified by immunogen affinity chromatography.

Specificity Recognizes endogenous levels of S100-A4 protein.

Clonality Monoclonal

Conjugation

Form Mouse IgG1. Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium

azide.

Dilution IH (1/100 - 1/300)

Gene Symbol S100A4

Alternative Names CAPL; MTS1; Protein S100-A4; Calvasculin; Metastasin; Placental calcium-binding

protein; Protein Mts1; S100 calcium-binding protein A4

Entrez Gene 6275 (Human)

SwissProt P26447 (Human)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

freeze/thaw cycles.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com










Immunohistochemical analysis of S100-A4 staining in human cutaneous squamous cell carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section



Immunohistochemical analysis of S100-A4 staining in human malignant melanoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish