

Product Data Sheet

Anti-CALCOCO1 Antibody

Catalog #	Source	Reactivity	Applications
CQA1710	Rabbit	H, M, R	WB, IH, IF/IC
Description	Rab	bit polyclonal antibody t	to CALCOCO1
Immunogen	Rec	ombinant full length pro	tein of human CALCOCO1
Purification	The	antibody was purified b	y immunogen affinity chromatography.
Specificity	Rec	ognizes endogenous leve	els of CALCOCO1 protein.
Clonality	Poly	clonal	
Conjugation			
Form	Liqu	uid in 0.42% Potassium p	hosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,
	and	0.01% sodium azide.	
Dilution	WB	(1/500 - 1/2000), IH (1/50	- 1/200), IF/IC (1/50 - 1/100)
Gene Symbol	CAL	.COCO1	
Alternative Na	ames KIA/	A1536; Calcium-binding	and coiled-coil domain-containing protein 1; Calphoglin;
	Coil	ed-coil coactivator prote	in; Sarcoma antigen NY-SAR-3
Entrez Gene	576	58 (Human); 67488 (Mo	use); 100912282, 246047 (Rat)
SwissProt	Q9P	P1Z2 (Human); Q8CGU1	(Mouse); Q66HR5 (Rat)
Storage/Stabi	lity Ship	oped at 4 $^\circ$ C. Upon deliv	very aliquot and store at -20 $^\circ$ C for one year. Avoid
	free	ze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



200


140


kDa A B C D

For research purposes only, not for human use

Product Data Sheet

Western blot analysis of CALCOCO1 expression in MCF7 (A), A549 (B), mouse lung (C), mouse heart (D) whole cell lysates. (Predicted band size: 67; 70; 77 kD; Observed band size: 77 kD)





Immunohistochemical analysis of CALCOCO1 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of CALCOCO1 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 $^{\circ}$ C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

 Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference
Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com