

Product Data Sheet

ADH5 siRNA (Human)

Source	Reactivity	Applications		
Synthetic	Н	RNAi		
siRNA	Գ to inhibit ADH5 expı	ession using RNA interference		
ADH5	ADH5 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
knocl	k down gene expressi	on.		
Lyoph	Lyophilized powder			
ADH5	ADH5			
mes ADH>	<; FDH; Alcohol dehyd	rogenase class-3; Alcohol dehydrogenase 5; Alcohol		
dehy	drogenase class chi ch	ain; Alcohol dehydrogenase class-III;		
Gluta	thione-dependent fo	maldehyde dehydrogenase; FALDH; FDH; GSH-FDH;		
S-(hy	droxymethyl)glutathio	one dehydrogenase		
128 (128 (Human)			
P117	P11766 (Human)			
> 97%	6			
l Oligo	nucleotide synthesis	s monitored base by base through trityl analysis to ensure		
appro	opriate coupling effici	ency. The oligo is subsequently purified by affinity-solid		
phase	e extraction. The anne	ealed RNA duplex is further analyzed by mass		
spect	rometry to verify the	exact composition of the duplex. Each lot is compared to		
the p	revious lot by mass sp	pectrometry to ensure maximum lot-to-lot consistency.		
We o	ffers pre-designed set	s of 3 different target-specific siRNA oligo duplexes of		
huma	an ADH5 gene. Each v	al contains 5 nmol of lyophilized siRNA. The duplexes can		
be tra	ansfected individually	or pooled together to achieve knockdown of the target		
gene,	, which is most comm	only assessed by qPCR or western blot.		
	Synthetic siRN/ ADHS knock Lyoph ADHS Mes ADHX dehy Gluta S-(hy 128 (P117 > 979 Oligo appro phase spect the p We o huma	Synthetic H siRNA to inhibit ADH5 expr ADH5 siRNA (Human) is a to knock down gene expression Lyophilized powder ADH5 mes ADH5 ADH5 mes ADHX; FDH; Alcohol dehyd dehydrogenase class chi chi Glutathione-dependent for S-(hydroxymethyl)glutathio 128 (Human) P11766 (Human) > 97% Oligonucleotide synthesis i appropriate coupling efficie phase extraction. The anne spectrometry to verify the the previous lot by mass sp We offers pre-designed set human ADH5 gene. Each vi be transfected individually		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Component	15 nmol	30 nmol
ADH5 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
ADH5 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
ADH5 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 µl
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com