

Product Data Sheet

CA3 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRH0529	Synthetic	н	RNAi		
Description	siRNA	siRNA to inhibit CA3 expression using RNA interference			
Specificity	CA3 s	CA3 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	knock down gene expression.			
Form	Lyoph	Lyophilized powder			
Gene Symbol	CA3	CA3			
Alternative N	ames Carbo	Carbonic anhydrase 3; Carbonate dehydratase III; Carbonic anhydrase III; CA-III			
Entrez Gene	761 (I	Human)			
SwissProt	P0745	P07451 (Human)			
Purity >		> 97%			
Quality Contr	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
appropriate coupling efficiency. The oligo is subsequently purifie			ified by affinity-solid		
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass sp	pectrometry to ensure maximum lo	t-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human CA3 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can			
	be tra	ansfected individually	or pooled together to achieve kno	ckdown of the target	
	gene, which is most commonly assessed by qPCR or western blot.			blot.	
	Com	ponent	15 nmol	30 nmol	
	CA3	siRNA (Human) - A	5 nmol x 1	5 nmol x 2	

CA3 siRNA (Human) - B 5 nmol x 1 5 nmol x 2 Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC Water	1 ml x 1	1 ml x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
CA3 siRNA (Human) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 μl
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com