

Product Data Sheet

IFNG siRNA (Human)

Catalog # Source Reactivity Applications

CRH2344 Synthetic H RNAi

Description siRNA to inhibit IFNG expression using RNA interference

Specificity IFNG siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to

knock down gene expression.

Form Lyophilized powder

Gene Symbol IFNG

Alternative Names Interferon gamma; IFN-gamma; Immune interferon

Entrez Gene 3458 (Human)

SwissProt P01579 (Human)

Purity > 97%

Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure

appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid

phase extraction. The annealed RNA duplex is further analyzed by mass

spectrometry to verify the exact composition of the duplex. Each lot is compared to

the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.

Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of

human IFNG gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can

be transfected individually or pooled together to achieve knockdown of the target

gene, which is most commonly assessed by qPCR or western blot.

| Component | 15 nmol | 30 nmol |
|------------------------|------------|------------|
| IFNG siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
| IFNG siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED



Product Data Sheet

| IFNG siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
|------------------------|--------------|--------------|
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μl of DEPC water to get a final concentration of 20 μM.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| 96-well | | 100 nM | 0.5 μΙ | 0.25 μΙ |
| | 100 μΙ | 50 nM | 0.25 μΙ | 0.25 μΙ |
| | | 10 nM | 0.05 μΙ | 0.25 μΙ |
| 24-well | | 100 nM | 2.5 μΙ | 1 μΙ |
| | 500 μl | 50 nM | 1.25 μΙ | 1 μΙ |
| | | 10 nM | 0.25 μΙ | 1 μΙ |
| 12-well | | 100 nM | 5 μΙ | 2 μΙ |
| | 1 ml | 50 nM | 2.5 μΙ | 2 μΙ |
| | | 10 nM | 0.5 μΙ | 2 μΙ |
| 6-well | | 100 nM | 10 μΙ | 5 μΙ |
| | 2 ml | 50 nM | 5 μΙ | 5 μΙ |
| | | 10 nM | 1 μΙ | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED