

# **Product Data Sheet**

### **IGFALS siRNA (Human)**

Catalog #	Source	Reactivity		Applications		
CRH2354	Synthetic	н		RNAi		
Description	siRNA	to inhibit IGFALS exp	pression using F	RNA interference		
Specificity	IGFAL	S siRNA (Human) is a	target-specific	19-23 nt siRNA oligo	duplexes designed to	
	knock	down gene expressi	on.			
Form	Lyoph	ilized powder				
Gene Symbol	IGFAL	IGFALS				
Alternative N	lames ALS; I	ALS; Insulin-like growth factor-binding protein complex acid labile subunit; ALS				
Entrez Gene	3483	3483 (Human)				
SwissProt	P3585	P35858 (Human)				
Purity	> 97%	> 97%				
Quality Conti	rol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	priate coupling effici	ency. The oligo	is subsequently puri	fied by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pi	revious lot by mass s	pectrometry to	ensure maximum lot	-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	human IGFALS gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can b	can be transfected individually or pooled together to achieve knockdown of the				
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent		15 nmol	30 nmol	
	IGFA	LS siRNA (Human) - A	A	5 nmol x 1	5 nmol x 2	
	IGFA	LS siRNA (Human) - E	3	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Negative Control2.5 nmol x 12.5 nmol x 2		5		
		DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
24-well		100 nM	2.5 μl	1 µl
	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
12-well		100 nM	5 μl	2 µl
	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com