

Product Data Sheet

SH3GL3 siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRH4367	Synthetic	Н	RNAi			
Description	siRNA	siRNA to inhibit SH3GL3 expression using RNA interference				
Specificity	SH3GL	SH3GL3 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expression	on.			
Form	Lyoph	ilized powder				
Gene Symbol	SH3GL	SH3GL3				
Alternative N	ames CNSA3	CNSA3; SH3D2C; Endophilin-A3; EEN-B2; Endophilin-3; SH3 domain protein 2C; SH3				
	domai	n-containing GRB2-li	ke protein 3			
Entrez Gene	6457 (Human)				
SwissProt	Q9996	Q99963 (Human)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligon	iucleotide synthesis i	s monitored base by base throu	gh trityl analysis to ensure		
	appro	priate coupling effici	ency. The oligo is subsequently p	ourified by affinity-solid		
	phase	extraction. The anne	ealed RNA duplex is further analy	yzed by mass		
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	evious lot by mass sp	pectrometry to ensure maximum	n lot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	humai	human SH3GL3 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can be	e transfected individu	ally or pooled together to achie	ve knockdown of the		
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
			A 5 nmol x 1	5 nmol x 2		
can be transfected individually or pooled together to achieve knockdown of the target gene, which is most commonly assessed by qPCR or western blot. Component 15 nmol 30 nmol			ve knockdown of the western blot. 30 nmol			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

SH3GL3 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
SH3GL3 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com