

Product Data Sheet

WHSC1 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|-----------|---|--|--|--|--|
| CRH5108 | Synthetic | н | RNAi | | | |
| Description | siRN | IA to inhibit WHSC1 e | xpression using RNA interference | | | |
| Specificity | WH | SC1 siRNA (Human) is | a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | kno | ck down gene expres | sion. | | | |
| Form | Lyop | philized powder | | | | |
| Gene Symbol | WH | SC1 | | | | |
| Alternative Na | ames KIAA | KIAA1090; MMSET; NSD2; TRX5; Histone-lysine N-methyltransferase NSD2; Multiple | | | | |
| | mye | loma SET domain-co | ntaining protein; MMSET; Nuclear SET domain-containing | | | |
| | prot | ein 2; NSD2; Protein | trithorax-5; Wolf-Hirschhorn syndrome candidate 1 protein; | | | |
| | WH | SC1 | | | | |
| Entrez Gene | 746 | 7468 (Human) | | | | |
| SwissProt | 096 | 028 (Human) | | | | |
| Purity | > 97 | 1% | | | | |
| Quality Contro | ol Olig | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | |
| | арр | ropriate coupling effi | ciency. The oligo is subsequently purified by affinity-solid | | | |
| | pha | se extraction. The an | nealed RNA duplex is further analyzed by mass | | | |
| | spee | ctrometry to verify th | e exact composition of the duplex. Each lot is compared to | | | |
| | the | previous lot by mass | spectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We | offers pre-designed s | ets of 3 different target-specific siRNA oligo duplexes of | | | |
| | hum | nan WHSC1 gene. Eac | h vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| | can | be transfected individ | dually or pooled together to achieve knockdown of the | | | |
| | targ | et gene, which is mos | st commonly assessed by qPCR or western blot. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|-------------------------|--------------|--------------|
| WHSC1 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
| WHSC1 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| WHSC1 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 μΙ |
| | | 50 nM | 5 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com