

Product Data Sheet

ZIC2 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---|--|---|---|------------------------|--|
| CRH5157 | Synthetic | н | RNAi | | |
| Description | siRNA | siRNA to inhibit ZIC2 expression using RNA interference | | | |
| Specificity | ZIC2 s | ZIC2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | knock down gene expression. | | | |
| Form | Lyoph | Lyophilized powder | | | |
| Gene Symbol | ZIC2 | ZIC2 | | | |
| Alternative N | ames Zinc fi | Zinc finger protein ZIC 2; Zinc finger protein of the cerebellum 2 | | | |
| Entrez Gene | 7546 | 7546 (Human) | | | |
| SwissProt | 09540 | O95409 (Human) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contr | ol Oligor | nucleotide synthesis is | nthesis is monitored base by base through trityl analysis to ensure | | |
| | appro | appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid | | | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | specti | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | |
| | the pr | revious lot by mass sp | ectrometry to ensure maximum lo | ot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | n ZIC2 gene. Each vial | contains 5 nmol of lyophilized siF | RNA. The duplexes can | |
| be transfected individually or pooled together to achieve knockdown o | | | ockdown of the target | | |
| | gene, which is most commonly assessed by qPCR or western blot. | | | blot. | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | ZIC2 | siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 | |

ZIC2 siRNA (Human) - B Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| | 1111 × 1 | 1111 × 2 | |
|------------------------|--------------|--------------|--|
| DEPC Water | 1 ml x 1 | 1 ml x 2 | |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 | |
| ZIC2 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 | |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 µl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 µl |
| | | 10 nM | 0.5 μl | 2 µl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 µl |
| | | 10 nM | 1 µl | 5 μl |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |