

Product Data Sheet

OGT siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|--|--|-----------------------------------|---------------------------|--|
| - | | | | | |
| CRH5555 | Synthetic | Н | RNAi | | |
| Description | Description siRNA to inhibit OGT expression using RNA interference | | | | |
| Specificity | OGT s | OGT siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | down gene expressio | n. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | OGT | OGT | | | |
| Alternative N | ames UDP-N | UDP-N-acetylglucosaminepeptide N-acetylglucosaminyltransferase 110 kDa | | | |
| | subur | nit; O-GlcNAc transfera | ase subunit p110; O-linked N-ace | tylglucosamine | |
| | transf | erase 110 kDa subuni | t; OGT | | |
| Entrez Gene | 8473 | (Human) | | | |
| SwissProt | 01529 | O15294 (Human) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contr | ol Oligor | Oligonucleotide synthesis is monitored base by base through trityl analysis to er | | | |
| | appro | priate coupling efficie | ncy. The oligo is subsequently pu | urified by affinity-solid | |
| | phase | extraction. The anne | aled RNA duplex is further analyz | zed by mass | |
| | specti | rometry to verify the e | exact composition of the duplex. | Each lot is compared to | |
| | the pr | revious lot by mass sp | ectrometry to ensure maximum | lot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | n OGT gene. Each vial | contains 5 nmol of lyophilized si | RNA. The duplexes can | |
| | be tra | nsfected individually | or pooled together to achieve kn | ockdown of the target | |
| | gene, | which is most commo | only assessed by qPCR or westerr | n blot. | |
| | Com | ponent | 15 nmol | 30 nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| OGT siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
|-----------------------|--------------|--------------|
| OGT siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| OGT siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com