

# **Product Data Sheet**

## **IKBKG siRNA (Human)**

| Catalog #      | Source     | Reactivity   | Applications  |  |  |
|----------------|------------|--|---|--|--|
| CRH5590        | Synthetic  | н  | RNAi  |  |  |
| Description    | siRNA      | to inhibit IKBKG exp   | ression using RNA interference                              |  |  |
| Specificity    | IKBKG      | siRNA (Human) is a   | target-specific 19-23 nt siRNA oligo duplexes designed to   |  |  |
|                | knock      | down gene expressi   | on.   |  |  |
| Form           | Lyoph      | nilized powder   |   |  |  |
| Gene Symbol    | IKBKG      | 5  |   |  |  |
| Alternative Na | ames FIP3; | FIP3; NEMO; NF-kappa-B essential modulator; NEMO; FIP-3; IkB kinase-associated |   |  |  |
|                | prote      | in 1; IKKAP1; Inhibito   | r of nuclear factor kappa-B kinase subunit gamma;           |  |  |
|                | I-kapp     | oa-B kinase subunit g  | amma; IKK-gamma; IKKG; IkB kinase subunit gamma;            |  |  |
|                | NF-ka      | ppa-B essential mod  |   |  |  |
| Entrez Gene    | 8517       | 8517 (Human)   |   |  |  |
| SwissProt      | Q9Y6       | K9 (Human)   |   |  |  |
| Purity         | > 97%      | Ď  |   |  |  |
| Quality Contro | ol Oligoi  | nucleotide synthesis   | is monitored base by base through trityl analysis to ensure |  |  |
|                | appro      | opriate coupling effici  | ency. The oligo is subsequently purified by affinity-solid  |  |  |
|                | phase      | e extraction. The anno   | ealed RNA duplex is further analyzed by mass                |  |  |
|                | spect      | rometry to verify the  | exact composition of the duplex. Each lot is compared to    |  |  |
|                | the p      | revious lot by mass s  | pectrometry to ensure maximum lot-to-lot consistency.       |  |  |
| Components     | We of      | ffers pre-designed se  | ts of 3 different target-specific siRNA oligo duplexes of   |  |  |
|                | huma       | in IKBKG gene. Each v  | ial contains 5 nmol of lyophilized siRNA. The duplexes can  |  |  |
|                | be tra     | ansfected individually   | or pooled together to achieve knockdown of the target       |  |  |
|                | gene,      | which is most comm   | only assessed by qPCR or western blot.                      |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| Component               | 15 nmol      | 30 nmol      |
|-------------------------|--------------|--------------|
| IKBKG siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
| IKBKG siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| IKBKG siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control        | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water              | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
| 96-well | 100 μl       | 100 nM              | 0.5 μl        | 0.25 μl      |
|         |              | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 µl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 µl         |
|         |              | 100 nM              | 5 µl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
| 6-well  | 2 ml         | 100 nM              | 10 µl         | 5 µl         |
|         |              | 50 nM               | 5 µl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com