

Product Data Sheet

PLA2G4C siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRH5653	Synthetic	Н	RNAi		
Description	siRNA	to inhibit PLA2G4C e	xpression using RNA interferen	се	
Specificity	PLA20	PLA2G4C siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expres	sion.		
Form	Lyoph	ilized powder			
Gene Symbol	PLA20	PLA2G4C			
Alternative Na	ames Cytoso	olic phospholipase A2	gamma; cPLA2-gamma; Phosp	holipase A2 group IVC	
Entrez Gene	8605 ((Human)			
SwissProt	Q9UP	65 (Human)			
Purity	> 97%				
Quality Contro	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling efficie	ency. The oligo is subsequently p	purified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	evious lot by mass sp	ectrometry to ensure maximun	n lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	n PLA2G4C gene. Eacl	h vial contains 5 nmol of lyophil	lized siRNA. The duplexes	
	can be	can be transfected individually or pooled together to achieve knockdown of the			
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	PLA2	G4C siRNA (Human) -	A 5 nmol x 1	5 nmol x 2	

PLA2G4C siRNA (Human) - B5 nmol x 15 nmol x 2Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-
Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility
Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
of medium	of siRNA		2000
	100 nM	0.5 μl	0.25 μl
100 µl	50 nM	0.25 μl	0.25 μl
	10 nM	0.05 μl	0.25 μl
	100 nM	2.5 μl	1 µl
500 μl	50 nM	1.25 μl	1 µl
	10 nM	0.25 μl	1 µl
	100 nM	5 µl	2 µl
1 ml	50 nM	2.5 μl	2 µl
	10 nM	0.5 μl	2 µl
2 ml	100 nM	10 µl	5 µl
	50 nM	5 µl	5 µl
	10 nM	1 µl	5 µl
	of medium 100 μl 500 μl 1 ml	of medium of siRNA 100 nM 100 nM 100 nM 10 nM 50 nM 10 nM 500 μl 50 nM 100 nM 10 nM 500 μl 50 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 1 nn 50 nM 10 nM 10 nM 10 nM 50 nM	100 nM0.5 μl100 μl50 nM0.25 μl10 nM0.05 μl100 nM2.5 μl500 μl50 nM1.25 μl10 nM0.25 μl10 nM50 μl100 nM5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com