

# **Product Data Sheet**

## ZBTB7B siRNA (Human)

| Catalog #       | Source    | Reactivity  | Applications  |  |  |
|-----------------|-----------|---|---|--|--|
| CRH9475         | Synthetic | Н   | RNAi  |  |  |
| Description     | siRN      | A to inhibit ZBTB7B ex  | pression using RNA interference                             |  |  |
| Specificity     | ZBTE      | 7B siRNA (Human) is   | a target-specific 19-23 nt siRNA oligo duplexes designed to |  |  |
|                 | knoc      | k down gene expressi  | on.   |  |  |
| Form            | Lyop      | hilized powder  |   |  |  |
| Gene Symbol     | ZBTE      | 57B   |   |  |  |
| Alternative Na  | ames ZBTE | 15; ZFP67; ZNF857B;   | Zinc finger and BTB domain-containing protein 7B;           |  |  |
|                 | Krue      | ppel-related zinc finge   | er protein cKrox; hcKrox; T-helper-inducing                 |  |  |
|                 | POZ/      | Krueppel-like factor;   | Zinc finger and BTB domain-containing protein 15; Zinc      |  |  |
|                 | finge     | er protein 67 homolog   | ; Zfp-67; Zinc finger protein 857B; Zinc finger protein     |  |  |
|                 | Th-P      | ОК  |   |  |  |
| Entrez Gene     | 5104      | 3 (Human)   |   |  |  |
| SwissProt       | 015:      | O15156 (Human)  |   |  |  |
| Purity          | > 97      | %   |   |  |  |
| Quality Control | ol Oligo  | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |   |  |  |
|                 | appr      | opriate coupling effici   | ency. The oligo is subsequently purified by affinity-solid  |  |  |
|                 | phas      | e extraction. The ann   | ealed RNA duplex is further analyzed by mass                |  |  |
|                 | spec      | trometry to verify the  | exact composition of the duplex. Each lot is compared to    |  |  |
|                 | the p     | previous lot by mass s  | pectrometry to ensure maximum lot-to-lot consistency.       |  |  |
| Components      | Wed       | offers pre-designed se  | ts of 3 different target-specific siRNA oligo duplexes of   |  |  |
|                 | hum       | an ZBTB7B gene. Each  | vial contains 5 nmol of lyophilized siRNA. The duplexes     |  |  |
|                 | can l     | pe transfected individ  | ally or pooled together to achieve knockdown of the         |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

target gene, which is most commonly assessed by qPCR or western blot.

| Component                | 15 nmol      | 30 nmol      |
|--------------------------|--------------|--------------|
| ZBTB7B siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
| ZBTB7B siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| ZBTB7B siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control         | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water               | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         | 100 µl       | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well |              | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
| 24-well | 500 μl       | 100 nM              | 2.5 μl        | 1 µl         |
|         |              | 50 nM               | 1.25 μl       | 1 μl         |
|         |              | 10 nM               | 0.25 μl       | 1 µl         |
| 12-well | 1 ml         | 100 nM              | 5 μl          | 2 μΙ         |
|         |              | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 µl         |
| 6-well  | 2 ml         | 100 nM              | 10 µl         | 5 μl         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

# **Product Data Sheet**

| 50 nM5 μl5 μl10 nM1 μl5 μl | o. /o. 1.1 |    |      |      |      |
|----------------------------|------------|----|------|------|------|
| 50 nM 5 μl 5 μl            |            | 10 | ) nM | 1 µl | 5 μl |
|                            |            | 50 | ) nM | 5 μl | 5 μΙ |

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com