

Product Data Sheet

TM6SF1 siRNA (Human)

siRNA to inhibit TM6SF1 expression using RNA interference			
esigned			
Transmembrane 6 superfamily member 1			
53346 (Human)			
Q9BZW5 (Human)			
to ensure			
appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid			
phase extraction. The annealed RNA duplex is further analyzed by mass			
spectrometry to verify the exact composition of the duplex. Each lot is compared to			
stency.			
kes of			
uplexes			
f the			
target gene, which is most commonly assessed by qPCR or western blot.			
; t s			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

TI	M6SF1 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Ν	egative Control	2.5 nmol x 1	2.5 nmol x 2
D	EPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
of medium	of siRNA		2000
	100 nM	0.5 μl	0.25 μl
100 µl	50 nM	0.25 μl	0.25 μl
	10 nM	0.05 μl	0.25 μl
	100 nM	2.5 μl	1 µl
500 μl	50 nM	1.25 μl	1 µl
	10 nM	0.25 μl	1 µl
	100 nM	5 µl	2 µl
1 ml	50 nM	2.5 μl	2 µl
	10 nM	0.5 μl	2 µl
	100 nM	10 µl	5 µl
2 ml	50 nM	5 µl	5 µl
	10 nM	1 µl	5 µl
	of medium 100 μl 500 μl 1 ml	of medium of siRNA 100 nM 100 nM 100 nM 10 nM 50 nM 10 nM 500 μl 50 nM 100 nM 10 nM 500 μl 50 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 1 nn 50 nM 10 nM 10 nM 10 nM 50 nM	100 nM0.5 μl100 μl50 nM0.25 μl10 nM0.05 μl100 nM2.5 μl500 μl50 nM1.25 μl10 nM0.25 μl10 nM50 μl100 nM5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com