

Product Data Sheet

Goat Anti-Quail IgY (H&L) - Biotin

Catalog #	Sourc	e Reactivity	Applications
CSA2278	Goat	Quail	E, WB, IH, IC
Description		Goat Polyclonal Secondary	Antibody to Quail IgY (H&L) Biotin labeled
Immunogen		Quail IgY	
Purification		The antibody was isolated	rom antisera by immunoaffinity chromatography using
		antigens coupled to agaros	e beads.
Specificity		By immunoelectrophoresis	and ELISA this antibody reacts specifically with Quail IgY.
		No antibody was detected	against non immunoglobulin serum proteins.
Clonality		Polyclonal	
Conjugation		Biotin	
Form		1 mg/ml. Liquid in 0.01M P	hosphate Buffered Saline, pH 7.2, containing 1% BSA,
		50% glycerol, 0.05% Sodiur	n Azide
Dilution		E (1/5000 - 1/20000), WB (1,	5000 - 1/20000), IH (1/500 - 1/1000), IC (1/500 - 1/1000)
Fluorophore			
Storage/Stabi	lity	Shipped at 4° C. Upon deli	very aliquot and store at -20 $^\circ$ C for one year. Avoid
		freeze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com