

Product Data Sheet

Goat Anti-Chicken IgG (H&L) - AMCA

Catalog #	Source	Reactivity	Applications
CSA2553	Goat	C	E, IH, IF, FC
Description	Goat Polyclonal Secondary Antibody to Chicken IgG (H&L) AMCA labeled		
Immunogen	Chicken IgG		
Purification	The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.		
Specificity	By immunoelectrophoresis and ELISA this antibody reacts specifically with Chicken IgG. No antibody was detected against non immunoglobulin serum proteins.		
Clonality	Polyclonal		
Conjugation	AMCA		
Form	1 mg/ml. Liquid in 0.01M Phosphate Buffered Saline, pH 7.2, containing 1% BSA, 50% glycerol, 0.02% Sodium Azide		
Dilution	IH (1/100 - 1/1000), IF (1/100 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay dependent concentration)		
Fluorophore	Ex/Em = 350/450 nm		
Storage/Stability	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com