

Product Data Sheet

Goat Anti-Human IgG+IgM+IgA - TRITC

Catalog #	Sourc	e Reactivity	Applications	
CSA2606	Goat	Н	E, IH, IF, FC	
Description		Goat Polyclonal Secondary Antibody to Human IgG+IgM+IgA TRITC labeled		
Immunogen		Human IgG+IgM+IgA		
Purification		The antibody was isolated from antisera by immunoaffinity chromatography using		
		antigens coupled to agaros	se beads.	
Specificity		By immunoelectrophoresis	and ELISA this antibody reacts specifically with Human	
		IgG IgM IgA. No antibody v	was detected against non immunoglobulin serum proteins.	
Clonality		Polyclonal		
Conjugation		TRITC		
Form		1 mg/ml. Liquid in 0.01M I	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,	
		50% glycerol, 0.02% Sodiu	m Azide	
Dilution		IH (1/100 - 1/1000), IF (1/10	0 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay	
		dependent concentration)		
Fluorophore		Ex/Em = 595/613 nm		
Storage/Stabi	ility	Shipped at 4°C. Upon deliv	ery aliquot and store at -20°C for one year. Avoid	
		freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com