

## **Product Data Sheet**

## Goat Anti-Human IgG Fc - Texas Red

Catalog #	Source	Reactivity	Applications		
CSA2704	Goat	н	E, IH, IF, FC		
Description		Goat Polyclonal Secondary Antibody to Human IgG Fc Texas Red labeled			
Immunogen		Human IgG Fc			
Purification		The antibody was isolated from antisera by immunoaffinity chromatography using			
		antigens coupled to agaros	e beads.		
Specificity		By immunoelectrophoresis and ELISA this antibody reacts specifically with Human			
		IgG Fc. No antibody was de	etected against non immunoglobulin serum proteins.		
Clonality		Polyclonal			
Conjugation		Texas Red			
Form		1 mg/ml. Liquid in 0.01M F	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,		
		50% glycerol, 0.02% Sodiu	n Azide		
Dilution		IH (1/100 - 1/1000), IF (1/10	0 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay		
		dependent concentration)			
Fluorophore		Ex/Em = 595/615 nm			
Storage/Stabi	lity	Shipped at 4°C. Upon deliv	ery aliquot and store at -20°C for one year. Avoid		
		freeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com