

Product Data Sheet

Goat Anti-Mouse IgM mu chain - Texas Red

Catalog #	Sourc	e Reactivity	Applications	
CSA2709	Goat	M	E, IH, IF, FC	
Description		Goat Polyclonal Secondary Antibody to Mouse IgM mu chain Texas Red labeled		
Immunogen		Mouse IgM mu chain		
Purification		The antibody was isolated from antisera by immunoaffinity chromatography using		
		antigens coupled to agard	ose beads.	
Specificity		By immunoelectrophores	sis and ELISA this antibody reacts specifically with Mouse	
		IgM mu chain. No antibo	dy was detected against non immunoglobulin serum	
		proteins.		
Clonality		Polyclonal		
Conjugation		Texas Red		
Form		1 mg/ml. Liquid in 0.01M	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,	
		50% glycerol, 0.02% Sodi	um Azide	
Dilution		IH (1/100 - 1/1000), IF (1/1	.00 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay	
		dependent concentration	n)	
Fluorophore		Ex/Em = 595/615 nm		
Storage/Stabi	ility	Shipped at 4°C. Upon del	ivery aliquot and store at -20°C for one year. Avoid	
		freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com