

Product Data Sheet

Mouse Anti-Duck IgG (H&L) - Texas Red

-				
Catalog #	Source	Reactivity	Applications	
CSA2757	Mouse	Dk	E, IH, IF, FC	
Description		Mouse Monoclonal Second	ary Antibody to Duck IgG (H&L) Texas Red labeled	
Immunogen		Duck IgG		
Purification		The antibody was isolated from ascitic by immunoaffinity chromatography using		
	i	antigens coupled to agarose	e beads.	
Specificity		By immunoelectrophoresis	and ELISA this antibody reacts specifically with Duck IgG.	
		No antibody was detected a	against non immunoglobulin serum proteins.	
Clonality		Monoclonal		
Conjugation		Texas Red		
Form		1 mg/ml. Liquid in 0.01M P	nosphate Buffered Saline, pH 7.2, containing 1% BSA,	
		50% glycerol, 0.02% Sodiun	n Azide	
Dilution		IH (1/100 - 1/1000), IF (1/100	- 1/1000), FC (1/1000 - 1/4000), E (Use at an assay	
		dependent concentration)		
Fluorophore		Ex/Em = 595⁄615 nm		
Storage/Stabi	lity	Shipped at 4°C. Upon delive	ry aliquot and store at -20°C for one year. Avoid	
	t	freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com